

ISO 22262-2:2014-09 (E)

Air quality - Bulk materials - Part 2: Quantitative determination of asbestos by gravimetric and microscopical methods

Contents		Page
Foreword		v
Introduction		vi
1	Scope	1
2	Normative references	1
3	Terms and definitions	1
4	Abbreviated terms	6
5	Determination of analytical requirements	6
6	Range	7
7	Limit of quantification	7
8	Principle	8
9	Safety precautions	8
10	Apparatus	8
11	Reagents	10
12	Sample size and homogeneity	10
12.1	Sample size	10
12.2	Representative sample	10
13	Methods for gravimetric matrix reduction	11
13.1	General	11
13.2	Data recording	11
13.3	Selection and pre-treatment of a representative sub-sample	13
13.4	Removal of organic materials by ashing	14
13.5	Acid treatment and sedimentation procedures	15
14	Procedures for quantification of asbestos in the final residue from gravimetric matrix reduction	18
14.1	General	18
14.2	Examination of the residue on the filter and selection of the appropriate procedure	19
15	Determination of asbestiform amphibole in vermiculite	25
15.1	General	25
15.2	Required size of sample for analysis	26
15.3	Sample pre-treatment	26
15.4	Separation of amphibole and measurement of the amphibole mass fraction	28
16	Determination of asbestos in talc	29
16.1	General	29
16.2	Determination of chrysotile in talc	29

16.3	Determination of amphibole in talc	29
17	Determination of compliance with legislative control limits	30
17.1	General	30
17.2	Gravimetry alone	30
17.3	Gravimetry combined with visual estimation	30
17.4	Gravimetry combined with point counting	30
17.5	Quantitative SEM or TEM fibre counting	32
18	Test report	32
Annex A (normative)	Types of commercial asbestos-containing materials and optimum analytical procedures	34
Annex B (normative)	Required centrifuge times for separation of amphibole in heavy liquid	42
Annex C (normative)	Example of test report	44
Bibliography		46