

ISO 22262-2:2026-01 (E)

Air quality - Bulk materials - Part 2: Quantitative determination of asbestos by gravimetric and microscopical methods

Contents		Page
Foreword		v
Introduction		vi
1	Scope	1
2	Normative references	1
3	Terms and definitions	1
4	Abbreviated terms	6
5	Determination of analytical requirements	6
6	Range	7
7	Limit of quantification	7
8	Principle	8
9	Safety precautions	8
10	Apparatus	8
11	Reagents	10
12	Sample size and homogeneity	10
12.1	Sample size	10
12.2	Representative sample	10
13	Methods for gravimetric matrix reduction	11
13.1	General	11
13.2	Data recording	11
13.3	Selection and pre-treatment of a representative sub-sample	13
13.3.1	General	13
13.3.2	Plasters without aggregate	13
13.3.3	Plasters with aggregate	13
13.3.4	Cements with and without aggregate	13
13.3.5	Floor tiles	13
13.3.6	Asphaltic materials without aggregate	14
13.3.7	Asphaltic materials with aggregate	14
13.3.8	Caulkings, mastics, putties, groutings and wall joint compounds	14
13.3.9	Cellulosic materials	14
13.3.10	Textured coatings	14
13.4	Removal of organic materials by ashing	14
13.4.1	General	14
13.4.2	Procedure	15
13.5	Acid treatment and sedimentation procedures	15
13.5.1	General	15
13.5.2	Procedure for acid treatment of samples containing soluble constituents, with or without insoluble aggregate	15

13.5.3	Procedure for resilient floor tile	17
13.5.4	Examination of materials for amphibole fibres	17
14	Procedures for quantification of the mass fraction of asbestos in the final residue from gravimetric matrix reduction	18
14.1	General	18
14.2	Examination of the residue on the filter and selection of the appropriate procedure	19
14.2.1	General	19
14.2.2	Gravimetric measurements alone	19
14.2.3	Visual estimation by PLM, SEM or TEM observation	19
14.2.4	Point counting by PLM or SEM	20
14.2.5	Determination of asbestos mass fraction from fibre measurements made by SEM or TEM	24
15	Determination of asbestos in talc and other mineral powders	26
15.1	General	26
15.2	Separation of chrysotile and amphibole by heavy liquid centrifugation	27
15.3	Determination of chrysotile and amphibole in talc and other mineral powders by TEM	27
15.3.1	General	27
15.3.2	Specimen preparation	27
15.3.3	Determination of chrysotile and amphibole mass fraction	28
15.3.4	Determination of chrysotile and amphibole numerical fibre concentrations	28
15.4	Determination of the mass fraction of chrysotile and amphibole in talc and other mineral powders by SEM	28
16	Determination of asbestiform amphibole in vermiculite	28
16.1	General	28
16.2	Required size of sample for analysis	29
16.3	Sample pre-treatment	29
16.3.1	Exfoliated vermiculite, loose fill vermiculite attic insulation and horticultural vermiculite	29
16.3.2	Vermiculite block fill insulation	30
16.3.3	Vermiculite ore concentrate	30
16.3.4	Vermiculite-containing horticultural products	30
16.3.5	Sprayed vermiculite-containing fireproofing	30
16.4	Separation of amphibole and measurement of the amphibole mass fraction	31
16.4.1	General	31
16.4.2	Manual separation of amphibole fibre bundles and weighing	31
16.4.3	Separation of amphibole by centrifugation in a heavy liquid	31
16.4.4	Confirmation of the absence of asbestos in the vermiculite sub-sample	32
17	Determination of compliance with legislative control limits	32
17.1	General	32
17.2	Gravimetry alone	32
17.3	Gravimetry combined with visual estimation	32
17.4	Gravimetry combined with point counting	33
17.5	Quantitative SEM or TEM fibre counting	35
18	Method validation	35
19	Test report	36
Annex A (normative)	Types of commercial asbestos-containing materials and optimum analytical procedures	37
Annex B (normative)	Required centrifuge times for separation of amphibole in heavy liquid	44
Annex C (normative)	Examples of test report	46
Bibliography	50