ISO 22262-2:2026-01 (E)

Air quality - Bulk materials - Part 2: Quantitative determination of asbestos by gravimetric and microscopical methods

Contents

Page

Foreword v

Introduction vi

1 Scope 1

2 Normative references 1

3 Termsanddefinitions 1

4 Abbreviated terms 6

5 Determination of analytical requirements 6

6 Range 7

7 Limitofquantification 7

8 Principle 8

9 Safety precautions 8

10 Apparatus 8

11 Reagents 10

12 Sample size and homogeneity 10
12.1 Sample size 10
12.2 Representative sample 10

13 Methods for gravimetric matrix reduction 11
13.1 General 11
13.2 Data recording 11
13.3 Selection and pre-treatment of a representative sub-sample 13
13.3.1 General 13
13.3.2 Plasters without aggregate 13
13.3.3 Plasters with aggregate 13
13.3.4 Cements with and without aggregate 13
13.3.5 Floor tiles 13
13.3.6 Asphaltic materials without aggregate 14
13.3.7 Asphaltic materials with aggregate 14
13.3.8 Caulkings, mastics, putties, groutings and wall joint compounds 14
13.3.9 Cellulosic materials 14
13.3.10 Textured coatings 14
13.4 Removal of organic materials by ashing 14
13.4.1 General 14
13.4.2 Procedure 15
13.5 Acid treatment and sedimentation procedures 15
13.5.1 General 15
13.5.2 Procedure for acid treatment of samples containing soluble constituents, with or without insoluble aggregate 15

13.5.3 Procedure for resilient floor tile 17
13.5.4 Examination of materials for amphibole fibres 17

14 Proceduresforquantificationofthemassfractionofasbestosinthefinalresiduefrom gravimetric matrix reduction 18
14.1 General 18
14.2 Examination of the residue on the filter and selection of the appropriate procedure 19
14.2.1 General 19
14.2.2 Gravimetric measurements alone 19
14.2.3 Visual estimation by PLM, SEM or TEM observation 19
14.2.4 Point counting by PLM or SEM 20
14.2.5 Determination of asbestos mass fraction from fibre measurements made by SEM or TEM 24

15 Determination of asbestos in talc and other mineral powders 26
15.1 General 26
15.2 Separation of chrysotile and amphibole by heavy liquid centrifugation 27
15.3 Determination of chrysotile and amphibole in talc and other mineral powders by TEM 27
15.3.1 General 27
15.3.2 Specimen preparation 27
15.3.3 Determination of chrysotile and amphibole mass fraction 28
15.3.4 Determination of chrysotile and amphibole numerical fibre concentrations 28
15.4 Determination of the mass fraction of chrysotile and amphibole in talc and other mineral powders by SEM 28

16 Determination of asbestiform amphibole in vermiculite 28
16.1 General 28
16.2 Required size of sample for analysis 29
16.3 Sample pre-treatment 29
16.3.1 Exfoliated vermiculite, loose fill vermiculite attic insulation and horticultural vermiculite 29
16.3.2 Vermiculite block fill insulation 30
16.3.3 Vermiculite ore concentrate 30
16.3.4 Vermiculite-containing horticultural products 30
16.3.5 Sprayed vermiculite-containing fireproofing 30
16.4 Separation of amphibole and measurement of the amphibole mass fraction 31
16.4.1 General 31
16.4.2 Manual separation of amphibole fibre bundles and weighing 31
16.4.3 Separation of amphibole by centrifugation in a heavy liquid 31
16.4.4 Confirmation of the absence of asbestos in the vermiculite sub-sample 32

17 Determination of compliance with legislative control limits 32
17.1 General 32
17.2 Gravimetry alone 32
17.3 Gravimetry combined with visual estimation 32
17.4 Gravimetry combined with point counting 33
17.5 Quantitative SEM or TEM fibre counting 35

18 Method validation 35

19 Test report 36

Annex A (normative) Types of commercial asbestos-containing materials and optimum analytical procedures 37

Annex B (normative) Required centrifuge times for separation of amphibole in heavy liquid 44

Annex C (normative) Examples of test report 46

Bibliography 50