

DIN EN 14476:2013-10 (E)

Chemical disinfectants and antiseptics - Quantitative suspension test for the evaluation of virucidal activity in the medical area - Test method and requirements (Phase 2/Step 1)

| Contents | | Page |
|--|---|-------------|
| Foreword | | 4 |
| Introduction | | 5 |
| 1 | Scope | 6 |
| 2 | Normative references | 6 |
| 3 | Terms and definitions | 6 |
| 4 | Requirements | 7 |
| 5 | Test methods | 8 |
| 5.1 | Principle | 8 |
| 5.2 | Materials and reagents, including cell cultures | 9 |
| 5.2.1 | Test organisms | 9 |
| 5.2.2 | Culture media, reagents and cell cultures | 9 |
| 5.3 | Apparatus and glassware | 12 |
| 5.4 | Preparation of test organism suspensions and product test solutions | 14 |
| 5.4.1 | Test organisms suspensions (test virus suspension) | 14 |
| 5.4.2 | Product test solutions | 14 |
| 5.5 | Procedure for assessing the virucidal activity of the product | 15 |
| 5.5.1 | General | 15 |
| 5.5.2 | Test procedure | 16 |
| 5.5.3 | Modified method for ready-to-use products | 17 |
| 5.5.4 | Cytotoxicity caused by product test solutions | 17 |
| 5.5.5 | Control of efficiency of suppression of product's activity | 18 |
| 5.5.6 | Reference test for virus inactivation | 18 |
| 5.5.7 | Titration of the virus control | 19 |
| 5.5.8 | Titration of test samples | 19 |
| 5.6 | Experimental data and calculation | 19 |
| 5.6.1 | Protocol of results | 19 |
| 5.6.2 | Calculation of infectivity titer (TCID ₅₀ or PFU) | 19 |
| 5.7 | Verification of the methodology | 19 |
| 5.8 | Expression of results | 20 |
| 5.8.1 | General | 20 |
| 5.8.2 | Calculation of the virucidal activity of products | 20 |
| 5.9 | Test report | 20 |
| Annex A (informative) Examples of viruses sorted according to their presence in the human body in case of virus infection | | 22 |
| Annex B (informative) Detoxification of test mixtures by molecular sieving | | 24 |
| B.1 | Molecular sieving with Sephadex TM LH 20 | 24 |
| B.1.1 | Principle | 24 |
| B.1.2 | Sephadex suspension | 24 |
| B.1.3 | Procedure | 24 |
| B.2 | Molecular sieving using MicroSpin TM S 400 HR | 26 |

| | |
|--|-----------|
| Annex C (informative) Calculation of the viral infectivity titre | 27 |
| C.1 Quantal tests Example of TCID₅₀ determination by the Spearman-Kärber method | 27 |
| C.2 Plaque test | 28 |
| C.3 Biometrical evaluation of experimental approaches and assessment of the disinfecting effect on the virus (reduction [R]): | 28 |
| C.3.1 General | 28 |
| C.3.2 Calculating the virus titre with 95 % confidence interval | 29 |
| C.3.3 Calculating the reduction and its 95 % confidence interval | 29 |
| C.3.4 Calculating the average reduction (R(mi)) and its 95 % confidence interval | 30 |
| C.3.5 Practical example | 31 |
| Annex D (informative) Presentation of test results of one active concentration | 33 |
| Annex E (informative) Quantitative determination of formaldehyde concentrations | 36 |
| Bibliography | 37 |