## ISO 20391-2:2019 (E)

# Biotechnology — Cell counting — Part 2: Experimental design and statistical analysis to quantify counting method performance

### Contents

Foreword

#### Introduction

- 1 Scope
- 2 Normative references
- 3 Terms, definitions, symbols and abbreviated terms
  - 3.1 Terms and definitions
    - 3.2 List of abbreviated terms and symbols
- 4 Principle
  - 4.1 General
  - 4.2 Proportionality
  - 4.3 Deviation from proportionality

#### 5 Experimental design

- 5.1 General
- 5.2 Considerations for the cell counting measurement process
- 5.3 Preparation of samples for the experimental design
- 5.3.1 General
- 5.3.2 Stock cell solution
- 5.3.3 Dilution fraction experimental design
- 5.3.4 Considerations for generating dilution fractions
- 5.4 Test sample labelling
- 5.5 Measurement of the test sample
- 6 Statistical methods
  - 6.1 General
  - 6.2 Mean cell count
  - 6.3 Measurement precision
  - 6.4 Proportional model fit
  - 6.5 Coefficient of determination
  - 6.6 Proportionality index (PI)
  - 6.6.1 General
  - 6.6.2 Calculation of the smoothed residual (esmoothed)
  - 6.6.3 Calculation of proportionality index (PI)
  - 6.7 Additional statistical analysis and quality metrics
  - 6.8 Data interpretation
  - 6.8.1 General
  - 6.8.2 Interpretation of %CV
  - 6.8.3 Interpretation of R2
  - 6.8.4 Interpretation of PI values
  - 6.8.5 Comparison of PI values
- 7 Reporting
  - 7.1 Reporting of quality indicators
  - 7.2 Documentation of experimental design parameters and statistical analysis method
  - 7.3 Additional reporting elements on the cell counting measurement process

#### Annex A (informative) Method to assess pipetting error contributions to dilution integrity

A.1 General

- A.2 Example procedure for pre-evaluating pipetting error contributions to dilution integrity using a calibrated scale to obtain accurate volume estimates upon pipetting
- A.3 Example procedure for obtaining a measured DF during the evaluation of PI using a calibrated scale to obtain accurate volume estimates
- Annex B (normative) Method to calculate smoothed residual (esmoothed) when a set of measured dilution fractions (DFmeasured) is obtained
- Annex C (informative) Example formulae for calculating PI
  - C.1 Detailed example of the calculation of PI based on sum of the absolute value of scaled smoothed residuals
  - C.2 Detailed example of the calculation of PI based on R2 of smoothed residuals
  - C.3 Additional examples of the calculation of PI when measured dilution fraction is utilized
- Annex D (informative) Use case 1 Evaluating the quality of a single cell counting measurement process
  - D.1 General
  - D.2 Description of experimental design and statistical analysis
  - D.3 Raw data and data analysis for use case 1
  - D.4 Example report for use case 1
  - D.5 Interpretation
- Annex E (informative) Use case 2 Comparing the quality of several cell counting measurement processes
  - E.1 General
  - E.2 Description of experimental design and statistical analysis
  - E.3 Raw data and data analysis for use case 2
  - E.4 Example report for use case 2
  - E.5 Comparison of cell counting methods evaluated in use case 2
  - E.5.1 Comparing quality indicators between cell counting methods
  - E.6 Interpretation
  - E.7 Consistency of results with underlying simulated model

Page count: 53