ISO 3517:2012-08 (E)

Essential oil of neroli (Citrus aurantium L., syn. Citrus amara Link, syn. Citrus bigaradia Loisel, syn. Citrus vulgaris Risso)

Contents

Page

Foreword iv

1 Scope 1

2 Normative references 1

3 Termsanddefinitions 1

4 Requirements 1
4.1 Appearance 1
4.2 Colour 2
4.3 Odour 2
4.4 Relative density at 20 °C, d20 20 2
4.5 Refractive index at 20 °C 2
4.6 Optical rotation at 20 °C 2
4.7 Miscibility with 85 % volume fraction ethanol at 20 °C 2
4.8 Acid value 2
4.9 Chromatographicprofile 2
4.10 Chirality 2
4.11 Flashpoint 2

5 Sampling 3

6 Test methods 3
6.1 Relative density at 20 °C, d20 20 3
6.2 Refractive index at 20 °C 3
6.3 Optical rotation at 20 °C 3
6.4 Miscibility in ethanol, 85 % volume fraction at 20 °C 3
6.5 Acid value 4
6.6 Chromatographicprofile 4
6.7 Chirality 4

7 Packaging, labelling and marking 4

Annex A (informative) Typical chromatograms of the analysis by gas chromatography of the essential oil of neroli (Citrus aurantium L., syn. Citrus amara Link, syn. Citrus bigaradia Loisel, syn. Citrus vulgaris Risso) 5

Annex B (informative) Flashpoint 8

Annex C (informative) Chirality 9

Bibliography 10