

ISO 19290:2016-12 (E)

Cigarettes - Determination of tobacco specific nitrosamines in mainstream cigarette smoke - Method using LC-MS/MS

Contents		Page
Foreword		iv
Introduction		v
1	Scope	1
2	Normative references	1
3	Terms and definitions	1
4	Principle	2
5	Apparatus	2
6	Reagents	2
7	Preparation	3
7.1	Preparation of glassware	3
7.2	Preparation of solutions	3
7.3	Preparation of standards	4
7.3.1	General	4
7.3.2	Preparation of internal standard solutions	4
7.3.3	Preparation of calibration standard solutions	4
8	Sampling	6
9	Tobacco product preparation	6
10	Sample generation -- Smoking of cigarettes	6
10.1	General	6
10.2	Linear smoking	6
10.3	Rotary smoking	6
11	Sample analysis	6
11.1	Preparation of sample	6
11.1.1	General	6
11.1.2	Extraction for linear smoking (44 mm pad)	6
11.1.3	Extraction for rotary smoking (92 mm pad)	6
11.1.4	Final sample preparation	6
11.2	Reversed phase high performance liquid chromatography	7
11.2.1	General	7
11.2.2	HPLC set-up parameters (Example)	7
11.2.3	Mobile phase (Example)	7
11.2.4	Mobile phase: Gradient (Example)	7
11.2.5	MS/MS Set-up parameters (Example)	7
11.3	Calculation	8
11.3.1	Calibration curve	8
11.3.2	Determination of the TSNA concentrations	9
11.3.3	Sample quantification	9
11.3.4	Determination of mainstream smoke TSNA deliveries	9

12	Repeatability and reproducibility	9
13	Test report	11
	Annex A (informative) Example of typical chromatograms	12
	Bibliography	14