

ISO 22634-2:2017-11 (E)

Cigarettes - Determination of benzo[a]pyrene in cigarette mainstream smoke using GC/MS - Part 2: Method using cyclohexane as extraction solvent

Contents		Page
Foreword		iv
Introduction		v
1	Scope	1
2	Normative references	1
3	Terms and definitions	1
4	Principle	1
5	Apparatus	2
6	Reagents	2
7	Standards	3
7.1	General	3
7.2	Primary B[a]P-d12 stock solution: 100 µg/ml	3
7.3	Secondary B[a]P-d12 spiking solution: 40 ng/ml	3
7.4	Primary B[a]P stock solution: 100 µg/ml	3
7.5	Secondary B[a]P stock solution: 1 000 ng/ml	3
7.6	Working standard solutions	3
7.7	Storage of standard solutions	3
8	Preparation of sample	3
8.1	Sampling	3
8.2	Smoking	4
8.3	Filter pad extraction	4
8.4	Sample clean-up	4
8.5	Blank solution	5
9	Determination	5
9.1	GC/MS operating conditions	5
9.2	Calibration	5
9.3	Determination of B[a]P	6
9.4	Calculation	6
10	Repeatability and reproducibility	6
11	Test report	7
11.1	General	7
11.2	Characteristic data about the cigarette	7
11.3	Data about sampling	7
11.4	Description of the test	8
11.5	Test results	8
Annex A (informative)	Example of a chromatogram of a cigarette smoke extract	9
Bibliography		10