

# ISO 11930:2019 (E)

## Cosmetics — Microbiology — Evaluation of the antimicrobial protection of a cosmetic product

---

### Contents

|           |  |
|-----------|--|
|           | Foreword   |
|           | Introduction   |
| 1         | Scope  |
| 2         | Normative references   |
| 3         | Terms and definitions  |
| 4         | Principle  |
| 5         | Preservation efficacy test   |
| 5.1       | General  |
| 5.2       | Materials, apparatus, reagents and culture media   |
| 5.2.1     | General  |
| 5.2.2     | Materials  |
| 5.2.3     | Diluents   |
| 5.2.3.1   | General  |
| 5.2.3.2   | Diluents for bacteria and <i>Candida albicans</i>  |
| 5.2.3.2.1 | Composition 1  |
| 5.2.3.2.2 | Preparation  |
| 5.2.3.2.3 | Composition 2  |
| 5.2.3.2.4 | Preparation  |
| 5.2.3.3   | Diluent for preparation of <i>Aspergillus brasiliensis</i> : polysorbate solution        |
| 5.2.4     | Neutralizer  |
| 5.2.4.1   | General  |
| 5.2.4.2   | Eugon LT 100 liquid broth  |
| 5.2.4.2.1 | General  |
| 5.2.4.2.2 | Composition  |
| 5.2.4.2.3 | Preparation  |
| 5.2.5     | Culture media  |
| 5.2.5.1   | General  |
| 5.2.5.2   | Culture medium for bacteria: tryptic soy agar (TSA) or soybean casein digest agar medium |
| 5.2.5.2.1 | Composition  |
| 5.2.5.2.2 | Preparation  |
| 5.2.5.3   | Culture medium for <i>C. albicans</i> : Sabouraud dextrose agar medium (SDA)             |
| 5.2.5.3.1 | Composition  |
| 5.2.5.3.2 | Preparation  |
| 5.2.5.4   | Culture medium for <i>A. brasiliensis</i> : potato dextrose agar (PDA)                   |
| 5.2.5.4.1 | Composition  |
| 5.2.5.4.2 | Preparation  |
| 5.3       | Microbial strains  |
| 5.4       | Preparation and enumeration of inocula   |
| 5.4.1     | General  |
| 5.4.2     | Preparation of bacterial and <i>Candida albicans</i> suspensions                         |
| 5.4.3     | Preparation of <i>Aspergillus brasiliensis</i> spore suspension                          |
| 5.5       | Demonstration of the neutralizer efficacy  |
| 5.5.1     | Principle  |
| 5.5.2     | Procedure  |
| 5.5.3     | Calculations   |
| 5.5.4     | Interpretation of results and conclusion on neutralizer efficacy                         |

- 5.6 Determination of the preservation efficacy of the formulation
  - 5.6.1 Procedure
    - 5.6.1.1 Sampling of test product
    - 5.6.1.2 Inoculation of test microorganisms
    - 5.6.1.3 Incubation of the inoculated formulation
    - 5.6.1.4 Sampling and enumeration
  - 5.6.2 Counting of colonies
  - 5.6.3 Calculations
    - 5.6.3.1 General
    - 5.6.3.2 Determination of the initial numbers of microorganisms, N and N0
    - 5.6.3.3 Enumeration of the microorganisms at each sampling time, Nx
    - 5.6.3.4 Reduction in microbial counts
  - 5.7 Interpretation of test results and conclusions
    - 5.7.1 Criteria
    - 5.7.2 General case (efficacy of the neutralizer is demonstrated for all strains)
    - 5.7.3 Case of formulations for which the efficacy of the neutralizer is not demonstrated for some strains
  - 5.8 Test report
- 6 Overall evaluation of the antimicrobial protection of the cosmetic product
  - 6.1 General
  - 6.2 Case 1 — Preservation efficacy test has been performed on the formulation
  - 6.3 Case 2 — Preservation efficacy test has not been performed on the formulation
- Annex A (normative) Decision diagram
- Annex B (normative) Evaluation criteria for the preservation efficacy test
- Annex C (informative) Examples of neutralizers for the antimicrobial activity of preservatives and washing liquids
- Annex D (informative) Packaging characteristics

Page count: 22