ISO 23906-1:2020 (E)

Cigarettes — Determination of benzo[a]pyrene in cigarette mainstream smoke with an intense smoking regime using GC/MS — Part 1: Method using methanol as extraction solvent

Contents

Foreword

Introduction

1 Scope

2 Normative references

3 Terms and definitions

4 Principle

5 Apparatus

6 Reagents

7 Standards

- 7.1 General
- 7.2 Primary B[a]P stock solution
- 7.3 Secondary B[a]P stock solution
- 7.4 B[a]P-d12 stock solution
- 7.5 B[a]P-d12 spiking solution
- 7.6 Working standard solutions
- 7.7 Storage of standard solutions

8 Preparation of sample

- 8.1 Sampling
- 8.2 Smoking
- 8.3 Glass-fibre filter pad extraction
- 8.4 Sample clean-up

9 Determination

- 9.1 GC-MS operating conditions
- 9.2 Calibration
- 9.3 Determination of B[a]P
- 9.4 Calculation

10 Repeatability and reproducibility

11 Test report

- 11.1 General
- 11.2 Characteristic data about the cigarette
- 11.3 Data about sampling
- 11.4 Description of the test
- 11.5 Test results

Annex A (informative) Example of a chromatogram of a cigarette smoke extract

Page count: 11