

# ISO 23906-2:2023-02 (E)

## Cigarettes - Determination of benzo[a]pyrene in cigarette mainstream smoke with an intense smoking regime using GC/MS - Part 2: Method using cyclohexane as extraction solvent

---

<b>Contents</b>		<b>Page</b>
Foreword .....		iv
Introduction .....		v
1	Scope .....	1
2	Normative references .....	1
3	Terms and definitions .....	1
4	Principles .....	1
5	Apparatus .....	2
6	Reagents .....	3
7	Standards .....	3
7.1	General .....	3
7.2	Primary B[a]P-d12 stock solution: 100 µg/ml .....	3
7.3	Secondary B[a]P-d12 spiking solution: 40 ng/ml .....	3
7.4	Primary B[a]P stock solution: 100 µg/ml .....	3
7.5	Secondary B[a]P stock solution: 1 000 ng/ml .....	3
7.6	Working standard solutions .....	3
7.7	Storage of standard solutions .....	3
8	Preparation of sample .....	4
8.1	Sampling .....	4
8.2	Smoking .....	4
8.3	Glass-fibre filter pad extraction .....	4
8.4	Sample clean-up .....	4
8.5	Blank solution .....	5
9	Determination .....	5
9.1	GC/MS operating conditions .....	5
9.2	Calibration .....	6
9.3	Determination of B[a]P .....	6
9.4	Calculation .....	6
10	Repeatability and reproducibility .....	6
11	Test report .....	7
11.1	General .....	7
11.2	Test results .....	7
Annex A (informative) Example of a chromatogram of a cigarette smoke extract .....		8
Bibliography .....		9