

DIN EN 14526:2026-06 (E)

Foodstuffs - Determination of saxitoxin-group toxins in shellfish - HPLC method using pre-column derivatization with peroxide or periodate oxidation

Contents		Page
Introduction		5
1	Scope	6
2	Normative references	7
3	Terms and definitions	7
4	Principle	7
5	Reagents	10
6	Apparatus	14
7	Procedure	15
7.1	Sample preparation	15
7.2	Extraction procedure	16
7.3	Sample purification	16
7.3.1	SPE-C18 clean-up	16
7.3.2	SPE-COOH clean-up (fractionation)	17
7.3.3	Alternative weak cation exchange SPE clean-up [2], [11]	17
7.4	Conversion of GTX6 into NEO and/or C3,4 into GTX1,4	18
7.4.1	General	18
7.4.2	Hydrolysis of SPE-COOH Fraction 1 or 2	18
7.5	Oxidation procedure	19
7.5.1	General	19
7.5.2	Periodate oxidation	19
7.5.3	Peroxide oxidation	20
8	HPLC determination	20
9	Calibration curve	23
10	Identification	23
11	Calculation	23
11.1	General	23
11.2	Calculation method with standard calibration curve	24
11.3	Calculation of GTX1,4 in the presence of GTX2,3 and dcGTX2,3	24
11.3.1	General	24
11.3.2	Method 1	26
11.4	Calculation of NEO in the presence of dcSTX	26
11.4.1	General	26
11.4.2	Method 2	28
11.5	Calculation of NEO in the presence of dcSTX, dcNEO and STX	28
11.5.1	General	28
11.5.2	Method 3	29
11.6	Calculation of dcNEO in the presence of dcSTX	30
11.6.1	General	30
11.6.2	Method 4	31
11.7	Calculation of C3,4	32

11.8	Calculation of GTX6	32
11.9	Conversion to STX 2HCl equivalents	32
12	Quality controls	33
12.1	General	33
12.2	Acceptance criteria series of analysis	33
12.3	Overall recovery	34
13	Verification	34
14	Test report	35
	Annex A (informative) Precision data	36
	Annex B (informative) Chromatograms	64
	Annex C (normative) Alternative calculation methods	68
	Bibliography	76