

ISO 13875:2005-02 (E)

Liquid milk - Determination of acid-soluble β-lactoglobulin content - Reverse-phase HPLC method

Contents		Page
Foreword		iv
1	Scope	1
2	Normative references	1
3	Terms and definitions	1
4	Principle	1
5	Reagents	1
6	Apparatus	2
7	Sampling	3
8	Procedure	3
8.1	Preparation of test portion	3
8.2	Preparation of test solution	4
8.3	Preparation of reference portion	4
8.4	Preparation of reference solutions for the multi-point calibration	4
8.5	Preparation of the reference solution for the "single-point" calibration procedure	5
8.6	HPLC determination	5
8.7	Integration mode	6
9	Calculation and expression of results	8
9.1	Multi-point calibration	8
9.2	Single-point calibration	9
9.3	Expression of results	9
10	Standardization of the reference sample	10
10.1	General	10
10.2	Preparation of the standard sample	10
10.3	Determination of protein content	10
10.4	Determination of -LG content in the reference sample	10
10.5	Calculation of the -LG content	10
10.6	Expression of results	11
11	Precision	11
11.1	Interlaboratory test	11
11.2	Repeatability	11
11.3	Reproducibility	11
12	Test report	12
Annex A (informative) Results of interlaboratory trials		13
Bibliography		14