ISO 18310-2:2021 (E)

Measurement and prediction of the ambient dose equivalent from patients receiving iodine 131 administration after thyroid ablation — Part 2: External effective dose of the caregivers after release from the hospital

Contents

Foreword

Introduction

1 Scope

2 Normative references

3 Terms and definitions

4 General requirements for the release of patient with caregiver

- 4.1 Discharge criteria
- 4.2 Management procedures of the patient receiving 131I administration
- 4.3 Release from the medical facility
- 4.4 Responsibility of the designated expert

5 Measurement of the effective dose to the caregiver

- 5.1 General
- 5.2 Specifications of the personal dosimeter
- 5.3 Measurement of the effective dose to the caregiver

6 Quality control

7 Uncertainty

Annex A (informative) Examples of written instructions to be presented to patients or their legal guardians before leaving the hospital after treatment with radioiodine

- A.1 Procedure for measuring radiation exposure to the caregiver of the patient receiving radiation treatment
- A.2 Recommended instructions for caregiver
- A.3 Precautions after discharge for patients and caregivers
- A.4 Letter of consent

Annex B (informative) Experimental application to this document: measurement and prediction of the effective dose to the caregiver in the vicinity of the patients receiving radioiodine 131 administration after thyroid ablation

- B.1 General
- B.2 Estimation of the effective dose to the caregiver
- B.3 Summary of measurement
- B.4 Result

Page count: 12