

DIN EN ISO 20031:2022-10 (E)

Radiological protection - Monitoring and dosimetry for internal exposures due to wound contamination with radionuclides (ISO 20031:2020)

Contents		Page
	European foreword	4
	Foreword	5
	Introduction	6
1	Scope	7
2	Normative references	7
3	Terms and definitions	7
4	Symbols and abbreviated terms	11
4.1	Symbols	11
4.2	Abbreviated terms	11
5	Purpose and need for special monitoring programmes for internal exposures due to wound contamination with radionuclides	11
6	General aspects of wound contamination	12
6.1	Introduction	12
6.2	Category of wound contaminants	12
6.3	Types of wounds and their specific retention of radionuclides	13
7	Monitoring programmes to assess contamination via a wound	13
7.1	Introduction	13
7.2	Main steps for the monitoring and dosimetry for internal exposures due to wound contamination with radionuclides	13
7.3	Collection of information to characterize the contaminated wound	14
7.3.1	General	14
7.3.2	Information concerning the type of wound	15
7.3.3	Information concerning the radioactive contaminant	15
7.4	In vivo wound measurements	15
7.5	Systemic activity monitoring	16
8	Performance criteria for radiobioassay measurements	17
9	Procedure for local and systemic dose assessment	17
9.1	Local (wound site) dose assessment	17
9.2	Systemic dose assessment	17
9.3	Impact of medical intervention on dose assessment	19
9.3.1	Local chelation therapy and/or the excision of contaminated tissue from the wound	19
9.3.2	Decorporation therapy	19
9.4	Software tools for bioassay data interpretation	19
9.5	Uncertainties	20
9.5.1	General	20
9.5.2	Uncertainties on local dose assessment	20
9.5.3	Uncertainties on internal dose assessment	20
9.6	Quality assurance	20
10	Recording	21
10.1	Recording in vivo measurement results	21
10.2	Recording in vitro radiobioassay and treatment waste results	21

11	Documentation of the dose assessment	22
12	Reporting	22
Annex A	(informative) Schematic representation of NCRP wound model, default parameters for retention equations and default transfer rates for the wound model for the various categories of radionuclides in wounds (adapted from NCRP report 156 (2007)^[3])	23
Annex B	(informative) Types of wounds and their specific retention of radionuclides	26
Annex C	(informative) Example of a summary sheet that should follow the contaminated worker during his initial care	28
Annex D	(informative) Overview of typical methods used for in vitro bioassay measurements	29
Annex E	(informative) Equivalent dose rate in a contaminated wound ($\text{mSv}\cdot\text{h}^{-1}\cdot\text{kBq}^{-1}$) and equivalent dose rate received by the skin ($\text{mSv}\cdot\text{h}^{-1}\cdot\text{kBq}^{-1}\cdot\text{cm}^2$) for selected radionuclides	30
Annex F	(informative) Committed effective dose coefficients for intake of selected radionuclides via a contaminated wound for all wound model categories (adapted from Toohey RE et al., 2014^[11])	32
Annex G	(informative) The IDEAS Guidelines^[14] provide guidelines for the estimation of committed doses from incorporation monitoring data in case of wound	35
Annex H	(informative) A-deviations	36
	Bibliography	37