

ISO 19040-1:2018 (E)

Water quality — Determination of the estrogenic potential of water and waste water — Part 1: Yeast estrogen screen (*Saccharomyces cerevisiae*)

Contents

	Foreword
1	Scope
2	Normative references
3	Terms and definitions
4	Principle
5	Interferences
6	Apparatus and materials
7	Reagents, media and test strain
8	Sampling and samples
8.1	General
8.2	Bottles and material for sampling
8.3	Bottles and material pre-cleaning
8.4	Sampling procedure
8.5	Transport of samples
8.6	Pretreatment of samples
8.7	Storage of samples
9	Procedure
9.1	Preparation of cryo-cultures for long-term storage
9.2	Overnight culture
9.3	Test set up for aqueous samples
9.3.1	Preparation
9.3.2	Preparation of the reference dilution series
9.3.3	Negative control
9.3.4	Blank replicate
9.3.5	Sample dilution
9.3.6	Field blank
9.3.7	Plate setup
9.3.8	Inoculation of the test plate
9.4	Measurement
9.4.1	Measurement of the cell density
9.4.2	Measurement of the reporter gene activity
9.5	Calculation of the corrected absorbance and the reporter gene induction
9.6	Calculation of the relative growth
9.7	Estimation of the EC50 of the reference compound by linear interpolation
10	Validity criteria
11	Assessment criteria
12	Test report
Annex A	(normative) Strain selection
A.1	General
A.2	Agar plates for strain selection

- A.3 Plating of the test strain
- Annex B (informative) Plate set up
- Annex C (informative) Scheme of test principle
- Annex D (informative) Test set up for chemicals and extracts
 - D.1 General
 - D.2 Extraction of water samples
 - D.3 Test with diluted organic solutions or extracts
 - D.4 Direct testing of organic solutions or extracts
 - D.5 Data from literature
- Annex E (informative) Preparation of dilution series
- Annex F (informative) Performance data
 - F.1 Design of the interlaboratory trial
 - F.1.1 General
 - F.1.2 Description of samples
 - F.1.3 Participating laboratories
 - F.2 Results of the interlaboratory trial
 - F.2.1 General
 - F.2.2 Summary of 17 β -estradiol equivalent concentrations (EEQ)
 - F.2.3 Summary of lowest ineffective dilutions LID
 - F.2.4 Trueness of results
- Annex G (informative) Use of other yeast strains based on *Saccharomyces cerevisiae*
 - G.1 General
 - G.2 Test strain according to Routledge and Sumpter [Reference (25)]
 - G.2.1 General
 - G.2.2 Strain description
 - G.2.3 Media
 - G.2.3.1 10x Su-DO-Medium (Sumpter).
 - G.2.3.2 Su-Growth medium (Sumpter).
 - G.2.3.3 Su-Exposure medium (Sumpter).
 - G.2.4 Strain selection
 - G.2.5 Procedure
 - G.2.6 Validity criteria
- Annex H (informative) Statistical assessment
- Annex I (informative) Calculation of 17 β -estradiol equivalents
 - I.1 General
 - I.2 Modelling of dose response relationship
 - I.3 Calculation of 17 β -estradiol equivalents for test samples
 - I.4 Reporting of estradiol equivalents for test samples
- Annex J (informative) Measurement of the lowest ineffective dilution (LID) of a waste water — A simplified evaluation for testing of waste water
 - J.1 General
 - J.2 Principle
 - J.3 Preparation of dilutions for LID assessment
 - J.4 Test for LID
 - J.5 Assessment of results — LID value, effluents
 - J.6 Documentation of results