ISO 19040-2:2018 (E)

Water quality — Determination of the estrogenic potential of water and waste water — Part 2: Yeast estrogen screen (A-YES, Arxula adeninivorans)

Contents

Foreword

- 1 Scope
- 2 Normative references
- 3 Terms and definitions
- 4 Principle
- 5 Interferences
- 6 Apparatus and materials
- 7 Reagents, media and test strains
- 8 Sampling and samples
 - 8.1 General
 - 8.2 Bottles and material for sampling
 - 8.3 Bottles and material pre-cleaning
 - 8.4 Sampling procedure
 - 8.5 Transport of samples
 - 8.6 Pretreatment of samples
 - 8.7 Storage of samples

9 Procedure

- 9.1 Test set up
- 9.1.1 Preparation of the reference dilution series
- 9.1.2 Reactivation of the yeast
- 9.1.3 Negative control
- 9.1.4 Blank replicate
- 9.1.5 Sample dilution
- 9.1.6 Field blank
- 9.1.7 Plate set up
- 9.1.8 Inoculation of the test plate
- 9.1.8.1 Preparation of test sample specific inoculation medium
- 9.1.8.2 Inoculation of wells without test sample
- 9.1.8.3 Inoculation of wells with test sample
- 9.2 Measurement
- 9.2.1 Measurement of the reporter gene activity
- 9.2.2 Measurement of the cell density
- 9.3 Calculations
- 9.3.1 Background correction
- 9.3.2 Calculation of the relative growth
- 9.3.3 Calculations for assessment of sample blanks
- 9.3.3.1 Calculations for assessment of sample coloration and artificial cleavage of substrate
- 9.3.3.2 Calculations for assessment of sample turbidity
- 9.3.4 Calculation of the reporter gene induction
- 10 Validity criteria
- 11 Assessment criteria

- 12 Test report
- 13 Verification
- Annex A (informative) Plate set up
 - A.1 General
 - A.2 Scheme of plate set up

Annex B (informative) Lyophilization of Arxula adeninivorans cells

- B.1 General
- B.2 Reagents
- B.3 Media
- B.3.1 Glucose solution
- B.3.2 Sorbitol solution
- B.3.3 PBS buffer
- B.3.4 Cryo medium
- B.3.5 Lysin solution
- B.3.6 Yeast minimal medium (YMM) with glucose
- B.3.7 Yeast minimal medium (YMM) with sorbitol
- B.4 Cultivation of Arxula adeninivorans on solid agar
- B.5 Cultivation of starter and main culture
- B.6 Preparing of lyophilized Arxula adeninivorans cells
- Annex C (informative) Scheme of test principle
- Annex D (informative) Test set up for chemicals and extracts
 - D.1 General
 - D.2 Extraction of water samples
 - D.3 Test with diluted organic solutions or extracts
- Annex E (informative) Preparation of dilution series
- Annex F (informative) Performance data
 - F.1 Validation study
 - F.1.1 Samples and participating laboratories
 - F.1.2 Results for lowest ineffective dilution (LID) and 17β-estradiol equivalent concentrations (EEQ)
 - F.1.3 Trueness evaluation
 - F.2 Sensitivity for selected compounds
- Annex G (informative) Statistical assessment
- Annex H (informative) Calculation of estradiol equivalents
 - H.1 General
 - H.2 Modelling of dose response relationship
 - H.3 Calculation of estradiol equivalents for test samples
 - H.4 Reporting of estradiol equivalents for test samples
- Annex I (informative) Alternative test design for EEQ determination
- Annex J (informative) Measurement of the lowest ineffective dilution (LID) of a waste water A simplified evaluation for testing of waste water
 - J.1 General
 - J.2 Principle
 - J.3 Preparation of dilutions for LID assessment
 - J.4 Test for LID
 - J.5 Assessment of results LID value, effluents
 - J.6 Documentation of results
- Annex K (informative) Example for statistical evaluation
 - K.1 General
 - K.2 Calculations
 - K.2.1 Background correction

- K.2.2 Calculation of the relative growth
- K.2.3 Calculations for assessment of sample blanks
- K.2.3.1 Sample coloration
- K.2.3.2 Artificial cleavage of substrate
- K.2.4 Calculations for assessment of sample turbidity
- K.2.5 Calculation of the report gene induction
- K.2.5.1 Calculation of the corrected absorbance for sample dilutions
- K.2.5.2 Calculation of the corrected absorbance for dilutions of the reference compound and negative control
- K.2.6 Calculation of induction rates

Page count: 55