

Soil quality — Determination of the effects of pollutants on soil flora — Leaf fatty acid composition of plants used to assess soil quality

Contents

	Foreword
	Introduction
1	Scope
2	Normative references
3	Terms, definitions and abbreviated terms
3.1	Terms and definitions
3.2	Abbreviated terms
4	Principle
5	Apparatus and reagents
5.1	Apparatus
5.2	Reagents
6	Sampling strategies
7	Sampling of leaf tissues
8	Obtaining, extraction and analyses of FAMES
8.1	Contamination control
8.2	Obtaining and extraction of FAMES from plant leaves
8.3	Analysis of FAMES
9	Test report
9.1	A reference to this document, i.e. ISO 21479
9.2	Description of the site and areas analysed
9.3	Leaf sampling
9.4	Fatty acid composition
9.5	Conclusion
Annex A	(informative) Results of the ring test
A.1	General
A.2	Results obtained in Bordeaux by six laboratories
A.3	Results obtained by the same investigator in different laboratories:
A.4	Outcome
Annex B	(informative) Assessment of soil quality by determining the Omega-3 index of <i>Lactuca sativa</i> seedlings grown ex situ under controlled conditions
Annex C	(informative) Plant species previously successfully used to assess soils of contaminated sites (organic and/or metals)
Annex D	(informative) Variation of the Omega-3 index as function of harvest time, plant size and leaf development
Annex E	(informative) Effect of the quantity of foliar tissues on the FAMES composition
Annex F	(informative) Example of chromatogram obtained after the FAMES analysis of foliar tissues
Annex G	(informative) Recommended mathematical method to rate soils of areas when some sampled plant species are not found in all areas