

ISO 18254-2:2018-12 (E)

Textiles - Method for the detection and determination of alkylphenol ethoxylates (APEO) - Part 2: Method using NPLC

Contents		Page
Foreword		iv
Introduction		v
1	Scope	1
2	Normative references	1
3	Terms and definitions	1
4	Principle	1
5	Reagents	1
6	Apparatus	2
7	Procedure	3
7.1	Standard preparation	3
7.2	Mobile phase for HPLC	3
7.2.1	Methanol with 0,1 % formic acid and 0,01 % ammonium formate	3
7.2.2	Acetonitrile with 0,1 % formic acid	3
7.3	Sample preparation	3
7.4	Sample extraction	3
7.5	Sample analysis	4
8	Calculation and calibration	4
8.1	Calculation of contribution percentage of each APEO congener in standard solution	4
8.2	Calculation of concentration of each APEO congener in standard solution	4
8.3	Calculation of concentration of each APEO congener in sample	4
8.4	Calculation of total concentration of all APEO congener in sample	5
8.5	Calibration curve	5
9	Test report	5
Annex A (informative)	Examples of chromatographic condition	6
Annex B (informative)	Examples of solid phase extraction (SPE) procedure for clean up	14
Annex C (informative)	Results of the interlaboratory test	15
Annex D (informative)	Study of the contribution percentage (mole fraction) of APEO congeners	16
Bibliography		18