

DIN EN ISO 18254-2:2019-09 (E)

Textiles - Method for the detection and determination of alkylphenol ethoxylates (APEO) - Part 2: Method using NPLC (ISO 18254-2:2018)

| Contents | Page |
|---|-------------|
| Figures Figure A.1 -- FLD chromatogram of NPEO using gradient condition (1) | 15 |
| Figure A.2 -- FLD chromatogram of OPEO using gradient condition (1) | 16 |
| Figure A.3 -- FLD chromatogram of NPEO using gradient condition (2) | 17 |
| Figure A.4 -- FLD chromatogram of OPEO using gradient condition (2) | 18 |
| Figure A.5 -- Overlay of FLD chromatograms of OPEO and NPEO using gradient condition (2) | 19 |
| Figure D.1 -- Comparison of contribution percentage | 23 |
| Tables Table A.1 -- Masses for the quantification of APEO in LC/MS | 13 |
| Table A.2 -- Masses for the quantification of APEO in LC/MS/MS | 14 |
| Table C.1 -- Statistical data of the interlaboratory test (sample with high concentration of APEO) | 21 |
| Table C.2 -- Statistical data of the interlaboratory test (sample with low concentration of APEO) | 21 |
| Table D.1 -- Average results of contribution percentage (mole fraction) | 22 |