

ISO 18218-2:2019 (E)

Leather — Determination of ethoxylated alkylphenols — Part 2: Indirect method

Contents

| | |
|---------|--|
| | Foreword |
| | Introduction |
| 1 | Scope |
| 2 | Normative references |
| 3 | Terms and definitions |
| 4 | Principle |
| 5 | Apparatus and materials |
| 6 | Chemicals |
| 7 | Sampling and sample preparation |
| 7.1 | Preparation of leather samples |
| 7.1.1 | Sampling and preparation of samples |
| 7.1.2 | Sample extraction |
| 7.2 | Preparation of leather process auxiliary samples |
| 7.3 | Blank determination |
| 7.4 | Determination of OP and NP |
| 7.5 | Determination of OPEO and NPEO |
| 7.6 | Chromatographic analysis |
| 7.7 | Evaluation |
| 8 | Calibration |
| 8.1 | Calibration for OP and NP |
| 8.2 | Calibration for OPEO and NPEO |
| 9 | Calculation |
| 9.1 | Calculation of OP and NP |
| 9.2 | Calculation of OPEO and NPEO |
| 10 | Test report |
| Annex A | (normative) Preparation of aluminium triiodide |
| A.1 | Reagents |
| A.2 | Apparatus |
| A.3 | Procedure |
| Annex B | (informative) Example of HPLC chromatograms |
| B.1 | HPLC conditions |
| Annex C | (informative) Example of GC-MS chromatograms |
| C.1 | Gas chromatographic conditions |
| C.2 | MS conditions |