

DIN EN ISO 20136:2020-12 (E)

Leather - Determination of degradability by micro-organisms (ISO 20136:2020)

| Contents | | Page |
|--|--|-------------|
| European foreword | | 3 |
| Foreword | | 4 |
| Introduction | | 5 |
| 1 | Scope | 6 |
| 2 | Normative references | 6 |
| 3 | Terms and definitions | 6 |
| 4 | Symbols and abbreviated terms | 7 |
| 5 | Principle | 7 |
| 5.1 | General | 7 |
| 5.2 | Assessment of biodegradation by manual titration; method A | 7 |
| 5.3 | Assessment of biodegradation by infrared (IR) detection; method B | 8 |
| 6 | Chemicals | 8 |
| 7 | Apparatus and materials | 9 |
| 8 | Procedure | 12 |
| 8.1 | Collection and preparation of the inoculum | 12 |
| 8.2 | Preparation of the test material and reference material | 12 |
| 8.3 | Test conditions and incubation period | 12 |
| 8.4 | Termination of the test | 12 |
| 9 | Quantification | 13 |
| 9.1 | Assessment of biodegradation by manual titration (method A) | 13 |
| 9.1.1 | Determination of the organic carbon content | 13 |
| 9.1.2 | Determination of the amount of CO₂ produced | 13 |
| 9.1.3 | Correcting for normality of HCl | 13 |
| 9.1.4 | Percentage of biodegradation from CO₂ evolved | 14 |
| 9.2 | Assessment of biodegradation by IR (method B) | 14 |
| 9.2.1 | Determination of the organic carbon content | 14 |
| 9.2.2 | Determination of the amount of CO₂ produced | 15 |
| 9.2.3 | Percentage of biodegradation from CO₂ data | 15 |
| 10 | Expression of results | 20 |
| 11 | Validity of results | 20 |
| 12 | Test report | 20 |
| Annex A (informative) Determination of the degree and rate of degradation of the material | | 21 |
| Annex B (informative) Quantitative determination of leather biodegradation | | 24 |
| Annex C (informative) Comparative biodegradability using different waste waters | | 28 |
| Bibliography | | 29 |