

DIN EN 15845:2010-05 (E)

Paper and board - Determination of the cytotoxicity of aqueous extracts

| Contents | | Page |
|-----------------|---|-------------|
| Foreword | | 3 |
| 1 | Scope | 4 |
| 2 | Normative references | 4 |
| 3 | Terms and definitions | 4 |
| 4 | Principle | 5 |
| 5 | Reagents | 5 |
| 6 | Cell line | 6 |
| 6.1 | Generating the cell strain | 6 |
| 6.2 | Maintaining the cell strain | 7 |
| 6.3 | Storing the cell strain | 7 |
| 7 | Food simulants used for testing | 7 |
| 8 | Cleaning laboratory glassware | 7 |
| 8.1 | Cleaning fluids for laboratory glassware | 7 |
| 8.2 | Cleaning procedure for laboratory glassware | 8 |
| 8.3 | Alternative cleaning procedure | 8 |
| 9 | Equipment | 8 |
| 9.1 | Equipment for the migration test | 8 |
| 9.2 | Cell culture equipment | 8 |
| 9.3 | Equipment used for cytotoxicity testing | 9 |
| 10 | Preparation of specimens | 10 |
| 10.1 | General | 10 |
| 10.2 | Paper and board intended for wet contact | 10 |
| 11 | Cytotoxicity assessment | 10 |
| 11.1 | Principle | 10 |
| 11.2 | General | 10 |
| 11.3 | Preparation of samples | 11 |
| 11.4 | Preparation of the reference sample, control sample and the positive control sample | 11 |
| 11.5 | Incubation of cells obtained from monolayer culture | 11 |
| 11.6 | Preparation of the chromatography sheet | 12 |
| 11.7 | Kinetics of uridine incorporation in the cell RNA | 12 |
| 12 | Expression of results | 13 |
| 12.1 | Graphic representation of the results (to be generated for each tube) | 13 |
| 12.2 | Calculation of percentage RNA synthesis and the validity of the test | 13 |
| 13 | Interpretation of the results | 15 |
| 13.1 | Results for the reference sample | 15 |
| 13.2 | Results of the positive control sample | 15 |
| 13.3 | Results for the test sample | 15 |
| 14 | Precision | 15 |

| | | |
|-----------|--|-----------|
| 15 | Test report | 16 |
| | Annex A (informative) Pyrodistilled water | 17 |
| | Bibliography | 18 |