ISO 20395:2019 (E)

Biotechnology — Requirements for evaluating the performance of quantification methods for nucleic acid target sequences — qPCR and dPCR

Contents

Foreword

Introduction

- 1 Scope
- 2 Normative references
- 3 Terms and definitions
- 4 Design of measurement procedure
 - 4.1 General
 - 4.2 Quantification method
 - 4.2.1 General
 - 4.2.2 qPCR determination of nucleic acid concentrations using a calibration curve
 - 4.2.3 dPCR determination of copy number concentration using molecular counting
 - 4.2.4 Relative quantification by qPCR
 - 4.2.5 dPCR determination of ratio between two targets
 - 4.3 Normalization strategy
 - 4.4 Controls
- 5 Sample QC Total nucleic acid quantity, integrity and purity
 - 5.1 General
 - 5.2 Total nucleic acid quantification
 - 5.2.1 General
 - 5.2.2 Spectrophotometry
 - 5.2.3 Fluorometry
 - 5.2.4 Assessment of total DNA concentration using qPCR/dPCR
 - 5.3 Nucleic acid integrity
 - 5.4 Nucleic acid purity
- 6 Assay design and optimization for quantification of nucleic acid target sequences
 - 6.1 Assay design
 - 6.1.1 General
 - 6.1.2 Amplicon selection
 - 6.1.3 Primer and probe design
 - 6.1.4 In silico evaluation of specificity
 - 6.1.5 RT-qPCR/RT-dPCR design
 - 6.2 Assay optimization using purified samples
 - 6.2.1 General
 - 6.2.2 Optimization of fluorescence signal
 - 6.2.3 (RT)-qPCR amplification efficiency
 - 6.2.4 RT efficiency
 - 6.2.5 Specificity
 - 6.3 Method optimization using test samples
 - 6.3.1 Effect of PCR inhibitors in sample matrix
 - 6.3.2 Presence of nucleic acid contaminants in test sample
 - 6.3.3 Validated measurement range
 - 6.4 No template controls

Data quality control (QC) and analysis

7.1 General

7

7.2 Acceptance criteria

- 7.2.1 qPCR
- 7.2.2 dPCR
- 7.3 Threshold setting
- 7.3.1 qPCR
- 7.3.2 dPCR
- 7.4 Data pre-processing
- 7.4.1 qPCR using calibration curve
- 7.4.2 Relative quantification (qPCR)
- 7.5 Identification of outliers
- Nucleic acid quantification measurement method validation
 - 8.1 General
 - 8.2 Precision
 - 8.3 LOQ
 - 8.4 LOD

8

- 8.5 Linearity
- 8.6 Trueness
- 8.7 Robustness
- 8.8 Specific considerations for qPCR method validation
- 8.8.1 Repeatability of qPCR- or RT-qPCR
- 8.8.2 Intermediate precision and reproducibility of qPCR- or RT-qPCR
- 8.9 Specific considerations for dPCR method validation
- 9 Nucleic acid quantification measurement traceability and comparability
 - 9.1 Metrological traceability
 - 9.2 Use of reference materials
 - 9.3 Instrument calibration
- 10 Measurement uncertainty (MU) in qPCR and dPCR measurements
 - 10.1 General requirements for MU calculations
 - 10.2 qPCR measurement uncertainty
 - 10.3 Ratio-based measurements
 - 10.4 dPCR measurement uncertainty
- 11 Reporting
- Annex A (informative) Spectrophotometry
 - A.1 General
 - A.2 Relationship between absorbance and total nucleic acid concentration
 - A.3 Absorbance ratios and nucleic acid purity

Annex B (informative) Nucleic acid integrity

- B.1 General
- B.2 Techniques for evaluation of DNA integrity
- B.2.1 Electrophoresis
- B.2.2 Long range PCR
- B.2.3 PCR assay with differential size amplicons
- B.3 Techniques for evaluation of RNA integrity
- B.3.1 Electrophoresis
- B.3.2 RT-qPCR for 5'/3' mRNA ratio
- B.3.3 RT-qPCR assay of repetitive elements
- B.3.4 RT-qPCR assays with different size amplicons
- Annex C (informative) PCR efficiency
 - C.1 General
 - C.2 Experimental design
 - C.3 Calculations
- Annex D (informative) Measurement uncertainty
 - D.1 General
 - D.2 Sources of uncertainty for whole process (qPCR)
 - D.3 Sources of uncertainty for dPCR
- Annex E (informative) MIQE and dMIQE checklists

- E.1 General
- E.2
- MIQE checklist dMIQE checklist E.3

Page count: 50