

ISO 17992:2013-09 (E)

Iron ores - Determination of arsenic content - Hydride generation atomic absorption spectrometric method

| Contents | | Page |
|-----------------------|---|-------------|
| Foreword | | iv |
| 1 | Scope | 1 |
| 2 | Normative references | 1 |
| 3 | Principle | 1 |
| 4 | Reagents | 1 |
| 5 | Apparatus | 3 |
| 6 | Sampling and samples | 4 |
| 6.1 | Laboratory sample | 4 |
| 6.2 | Preparation of predried test samples | 4 |
| 7 | Procedure | 4 |
| 7.1 | Number of determinations | 4 |
| 7.2 | Test portion | 4 |
| 7.3 | Blank test and check test | 4 |
| 7.4 | Determination | 5 |
| 8 | Expression of results | 7 |
| 8.1 | Calculation of mass fraction of arsenic content | 7 |
| 8.2 | General treatment of results | 7 |
| 8.3 | Oxide factor | 10 |
| 9 | Test report | 10 |
| Annex A (normative) | Flow sheet of the procedure for the acceptance of analytical values for test sample | 13 |
| Annex B (informative) | Parameters of some instruments | 14 |
| Annex C (informative) | Derivation of repeatability and permissible tolerance formulae | 16 |
| Annex D (informative) | Precision data obtained by international analytical trials | 17 |