ISO 21257:2018 (E)

Plastics — Polymer polyols for use in the production of polyurethane — Determination of the residual acrylonitrile and styrene monomer content by gas chromatography

Contents

Foreword

Introduction

1 Scope

2 Normative references

3 Terms and definitions

4 Principle

5 Reagents and materials

6 Apparatus

7 Preparation of calibration solution and sample solution

- 7.1 General
- 7.2 Method A for using OT column
- 7.2.1 Preparation of internal standard solution
- 7.2.2 Preparation of calibration solutions
- 7.2.3 Preparation of sample solution
- 7.3 Method B for using packed column
- 7.3.1 Preparation of internal standard solution
- 7.3.2 Preparation of calibration solutions
- 7.3.3 Preparation of sample solution

8 Procedure

- 8.1 Gas chromatographic procedure
- 8.2 Measurement of sample solutions and calibration solutions
- 8.3 Evaluation of gas chromatographic peak

9 Expression of the results

- 9.1 Preparation of the calibration curve
- 9.2 Calculation of results from a calibration graph
- 9.3 Calculation of results from a single-point calibration
- 9.4 Acceptability of results and measurement sensitivity

10 Precision

11 Test report

Annex A (informative) Typical analytical conditions

- A.1 Condition of chromatograph with OT column
- A.2 Condition of chromatograph with packed column

Annex B (informative) Typical examples of GC chromatogram

Annex C (informative) Precision

Page count: 15