

ISO 21257:2018 (E)

Plastics — Polymer polyols for use in the production of polyurethane — Determination of the residual acrylonitrile and styrene monomer content by gas chromatography

Contents

	Foreword
	Introduction
1	Scope
2	Normative references
3	Terms and definitions
4	Principle
5	Reagents and materials
6	Apparatus
7	Preparation of calibration solution and sample solution
7.1	General
7.2	Method A for using OT column
7.2.1	Preparation of internal standard solution
7.2.2	Preparation of calibration solutions
7.2.3	Preparation of sample solution
7.3	Method B for using packed column
7.3.1	Preparation of internal standard solution
7.3.2	Preparation of calibration solutions
7.3.3	Preparation of sample solution
8	Procedure
8.1	Gas chromatographic procedure
8.2	Measurement of sample solutions and calibration solutions
8.3	Evaluation of gas chromatographic peak
9	Expression of the results
9.1	Preparation of the calibration curve
9.2	Calculation of results from a calibration graph
9.3	Calculation of results from a single-point calibration
9.4	Acceptability of results and measurement sensitivity
10	Precision
11	Test report
Annex A	(informative) Typical analytical conditions
A.1	Condition of chromatograph with OT column
A.2	Condition of chromatograph with packed column
Annex B	(informative) Typical examples of GC chromatogram
Annex C	(informative) Precision