

# ISO 24076:2021 (E)

## Plastics — Polypropylene (PP) — Determination of isotactic index by low-resolution nuclear magnetic resonance spectrometry

---

### Contents

|         |   |
|---------|---|
|         | Foreword  |
|         | Introduction  |
| 1       | Scope   |
| 2       | Normative references                                      |
| 3       | Terms and definitions                                     |
| 4       | Principle   |
| 5       | Apparatus   |
| 6       | Preparation of sample                                     |
| 6.1     | General   |
| 6.2     | Reference samples   |
| 7       | Method A — Ratio method                                   |
| 7.1     | Procedure   |
| 7.1.1   | Measurement of reference samples by n-heptane extraction  |
| 7.1.2   | LR-NMR measurement procedure of reference samples         |
| 7.1.3   | LR-NMR measurement procedure of test samples              |
| 7.2     | Calculation   |
| 7.2.1   | Calibration curve   |
| 7.2.2   | Calculation of isotactic index $II_{na}$                  |
| 8       | Method B — Absolute method                                |
| 8.1     | Procedure   |
| 8.1.1   | Measurement of reference samples by n-heptane extraction  |
| 8.1.2   | LR-NMR measurement procedure of reference samples         |
| 8.1.3   | LR-NMR measurement procedure of test samples              |
| 8.2     | Calculation   |
| 8.2.1   | Calibration curve   |
| 8.2.2   | Calculation of isotactic index $II_{nb}$                  |
| 9       | Expression of results                                     |
| 10      | Test report   |
| Annex A | (informative) Example of a calibration curve for method A |
| Annex B | (informative) Example of a calibration curve for method B |