ISO/TS 16782:2016-10 (E)

Clinical laboratory testing - Criteria for acceptable lots of dehydrated Mueller-Hinton agar and broth for antimicrobial susceptibility testing

Contents

Page

Foreword iv

Introduction v

1 Scope 1

2 Normative reference 1

3 Termsanddefinitions 1

4 Requirements for Mueller-Hinton broth 3
4.1 Components of Mueller-Hinton broth 3
4.2 Physical and chemical characteristics 3
4.2.1 Dehydrated powder or granules 3
4.2.2 Prepared broth medium 4
4.2.3 Cation supplementation and content for MHB 4
4.2.4 Other medium components 4
4.2.5 Specific adjustments required by the manufacturer 4
4.3 Manufacturers protocol for testing production lots of dehydrated Mueller-Hinton broth 5
4.4 Interpreting the results 5
4.5 Evaluating the results 6

5 Requirements for Muller-Hinton agar 6
5.1 Components of Mueller-Hinton agar 6
5.2 Physical and chemical characteristics 6
5.2.1 Dehydrated powder or granules 6
5.2.2 Prepared agar medium 7
5.2.3 Cation supplementation and content for MHA 7
5.2.4 Other medium components 7
5.2.5 Specific adjustments required by the manufacturer 7
5.3 Manufacturer's protocol for testing production lots of dehydrated Mueller-Hinton agar 8
5.4 Interpreting the results 8
5.5 Evaluating the results 10

6 Testing new antimicrobial agents with production lots of dehydrated Mueller- Hinton broth or agar 11

Annex A (informative) Mueller-Hinton medium 12

Annex B (informative) Preparing control cultures 14

Annex C (informative) Suggested data sheet for testing of production lots 16

Annex D (informative) Label statement 19

Bibliography 20