DIN EN 16159:2012-04 (E)

Animal feeding stuffs - Determination of selenium by hydride generation atomic absorption spectrometry (HGAAS) after microwave digestion (digestion with 65 % nitric acid and 30 % hydrogen peroxide)

Contents

Page

Foreword 3

1 Scope 4

2 Normative references 4

3 Principle 4

4 Reagents 5

5 Apparatus and equipment 6

6 Procedure 7
6.1 General 7
6.2 Preparation of the test solution 7
6.3 Measurement of the test solution 8
6.3.1 Pre-dilution of the test solution 8
6.3.2 Pre-reduction of the (pre-diluted) test solution 8
6.3.3 Spectrometer settings of the flow-injection hydride generation atomic absorption spectrometer (HGAAS) 9
6.3.4 HGAAS determination 9

7 Calculation 10

8 Precision 10
8.1 Introduction 10
8.2 General 11
8.3 Repeatability 11
8.4 Reproducibility 11

9 Test report 11

Annex A (informative) Results of the inter-laboratory test 12

Annex B (informative) Flowchart - Determination of selenium by hydride generation atomic absorption spectrometry (HGAAS) after microwave digestion (digestion with 65 % nitric acid and 30 % hydrogen peroxide 13

Annex C (informative) Alternative digestion procedure with the same digestion efficiency: Acid digestion with a mixture of 65 % nitric acid and 70 % perchloric acid (7:3 by volume) at atmospheric pressure 14

C.1 Preparation of selenium calibration solutions 14
C.2 Preparation of the test solution 14
C.3 Measurement of the test solution 15
C.4 Ten rules for automated wet ashing with perchloric acid 15

Bibliography 16