

ISO 3517:2012-08 (E)

Essential oil of neroli (*Citrus aurantium* L., syn. *Citrus amara* Link, syn. *Citrus bigaradia* Loisel, syn. *Citrus vulgaris* Risso)

Contents		Page
Foreword		iv
1	Scope	1
2	Normative references	1
3	Terms and definitions	1
4	Requirements	1
4.1	Appearance	1
4.2	Colour	2
4.3	Odour	2
4.4	Relative density at 20 °C, d ₂₀ 20	2
4.5	Refractive index at 20 °C	2
4.6	Optical rotation at 20 °C	2
4.7	Miscibility with 85 % volume fraction ethanol at 20 °C	2
4.8	Acid value	2
4.9	Chromatographic profile	2
4.10	Chirality	2
4.11	Flashpoint	2
5	Sampling	3
6	Test methods	3
6.1	Relative density at 20 °C, d ₂₀ 20	3
6.2	Refractive index at 20 °C	3
6.3	Optical rotation at 20 °C	3
6.4	Miscibility in ethanol, 85 % volume fraction at 20 °C	3
6.5	Acid value	4
6.6	Chromatographic profile	4
6.7	Chirality	4
7	Packaging, labelling and marking	4
Annex A (informative)	Typical chromatograms of the analysis by gas chromatography of the essential oil of neroli (<i>Citrus aurantium</i> L., syn. <i>Citrus amara</i> Link, syn. <i>Citrus bigaradia</i> Loisel, syn. <i>Citrus vulgaris</i> Risso)	5
Annex B (informative)	Flashpoint	8
Annex C (informative)	Chirality	9
Bibliography		10