

ISO/TS 13136:2012-11 (E)

Microbiology of food and animal feed - Real-time polymerase chain reaction (PCR)-based method for the detection of food-borne pathogens - Horizontal method for the detection of Shiga toxin-producing Escherichia coli (STEC) and the determination of O157, O111, O26, O103 and O145 serogroups

Contents		Page
Foreword		iv
Introduction		v
1	Scope	1
2	Normative references	1
3	Terms and definitions	2
4	Principle	2
4.1	General	2
4.2	Microbial enrichment	2
4.3	Nucleic acid extraction	3
4.4	Target genes	3
4.5	Detection	3
4.6	Isolation	3
5	Diluents, culture media and reagents	3
5.1	Culture media	4
5.2	Reagents for nucleic acid extraction	5
5.3	Reagents for PCR	5
6	Equipment	5
7	Sampling	6
8	Preparation of test sample	6
9	Procedure	6
9.1	Test portion and initial suspension	6
9.2	Enrichment	7
9.3	Nucleic acid extraction	7
9.4	PCR amplification (for real-time PCR)	7
9.5	Strain isolation	8
10	Expression of results	8
11	Performance data	8
Annex A (normative)	Flow diagram of the screening procedure	12
Annex B (normative)	Flow diagram of the isolation and confirmation procedure	13
Annex C (informative)	Identification of Shiga toxin-producing Escherichia coli (STEC) by multiplex PCR amplification of virulence genes and detection of PCR products by agarose gel electrophoresis	14

Annex D (informative) Internal amplification control	18
Annex E (informative) Primers and probes for the PCR assays	19
Annex F (normative) Isolation of STEC strains	21
Bibliography	22