ISO 21330:2018 (E)

Cigarettes — Determination of selected volatile organic compounds in the mainstream smoke of cigarettes — Method using GC/MS

Contents

Foreword

Introduction

1 Scope

2 Normative references

3 Terms and definitions

4 Principle

5 Apparatus

6 Reagents

7 Preparation

7.1 Preparation of glassware
7.2 Preparation of standards
7.2.1 General
7.2.2 Preparation of internal standard spiking solution
7.2.2.1 Internal standard stock solution
7.2.2.2 Internal standard spiking solution
7.2.2.3 Storage
7.2.3 Preparation of working standards for isoprene, acrylonitrile, benzene, and toluene
7.2.3.1 Primary isoprene, acrylonitrile, benzene and toluene stock solutions
7.2.3.2 Secondary stock solution (mixture of isoprene, acrylonitrile, benzene and toluene primary stock solutions)
7.2.3.3 Calibration standard solutions (for isoprene, acrylonitrile, benzene and toluene)
7.2.3.4 Storage
7.2.4 Preparation of working standards for 1,3-butadiene
7.2.4.1 Primary 1,3-butadiene stock solution
7.2.4.2 Secondary 1,3-butadiene stock solution
7.2.4.3 Determination of secondary 1,3-butadiene stock concentration
7.2.4.4 1,3-butadiene calibration standard solutions
7.2.4.5 Storage

8 Sampling

9 Tobacco product preparation

10 Sample generation — Smoking of cigarettes

10.1 General
10.2 Smoking machine setup
10.3 Smoking
10.3.1 General
10.3.2 Linear smoking
10.3.3 Rotary smoking

11 Sample analysis

11.1 Preparation of sample
11.2 Determination
11.2.1 GC-MS operating conditions

11.2.2 Calibration
11.2.3 Calculation

12 Repeatability and reproducibility

12.1 General
12.2 Results of the 2012 collaborative study

13 Test report

Annex A (informative) Examples of chromatograms

Page count: 17