ISO 15216-2:2019 (E)

Microbiology of the food chain — Horizontal method for determination of hepatitis A virus and norovirus using real-time RT-PCR — Part 2: Method for detection

Contents

Foreword

Introduction

1 Scope

2 Normative references

3 Terms and definitions

4 Principle

4.1 Virus extraction
4.2 RNA extraction
4.3 Real-time RT-PCR
4.4 Control materials
4.4.1 Process control virus
4.4.2 EC RNA control
4.5 Test results

5 Reagents

5.1 General
5.2 Reagents used as supplied
5.3 Reagents requiring preparation

6 Equipment and consumables

7 Sampling

8 Procedure

8.1 General laboratory requirements
8.2 Virus extraction
8.2.1 General
8.2.2 Process control virus material
8.2.3 Negative process control
8.2.4 Surfaces
8.2.5 Soft fruit and leaf, stem and bulb vegetables
8.2.6 Bottled water
8.2.7 Bivalve molluscan shellfish (BMS)
8.3 RNA extraction
8.4 Real-time RT-PCR
8.4.1 General requirements
8.4.2 Real-time RT-PCR analysis
8.4.2.1 Analysis of target virus
8.4.2.2 Analysis of process control virus
8.4.2.3 Amplification
8.4.2.4 Analysis of fluorescence data

9 Interpretation of results

9.1 General
9.2 Construction of process control virus RNA standard curve
9.3 Control for RT-PCR inhibition
9.4 Calculation of extraction efficiency

10 Expression of results

11 Performance characteristics of the method

11.1 Validation study
11.2 Sensitivity
11.3 Specificity
11.4 LOD50

12 Test report

Annex A (normative) Diagram of procedure

Annex B (normative) Composition and preparation of reagents and buffers

B.1 5 × PEG/NaCl solution (500 g/l PEG 8 000, 1,5 mol/l NaCl)
B.1.1 Composition
B.1.2 Preparation
B.2 Chloroform/butanol mixture (1:1 volume fraction)
B.2.1 Composition
B.2.2 Preparation
B.3 Proteinase K solution (3000 U/l)
B.3.1 Composition
B.3.2 Preparation
B.4 Phosphate-buffered saline (PBS)
B.4.1 Composition
B.4.2 Preparation
B.5 Tris/glycine/beef extract (TGBE) buffer
B.5.1 Composition
B.5.2 Preparation
B.6 Tris solution (1 mol/l)
B.6.1 Composition
B.6.2 Preparation
B.7 EDTA solution (0,5 mol/l)
B.7.1 Composition
B.7.2 Preparation
B.8 Tris EDTA (TE) buffer (10 mmol/l Tris, 1 mmol/l EDTA)
B.8.1 Composition
B.8.2 Preparation

Annex C (informative) Real-time RT-PCR mastermixes and cycling parameters

Annex D (informative) Real-time RT-PCR primers and hydrolysis probes for the detection of HAV, norovirus GI and GII and mengo virus (process control)

D.1 HAV
D.2 Norovirus GI
D.3 Norovirus GII
D.4 Mengo virus

Annex E (informative) Growth of mengo virus strain MC0 for use as a process control

E.1 General
E.2 Reagents and apparatus
E.3 Procedure

Annex F (informative) RNA extraction using the BioMerieux NucliSens®3 3 BioMerieux NucliSens® is the trade name of a product supplied by BioMerieux. This information is given for the convenience of users of this document and does not constitute an endorsement by ISO of the product named. Equivalent products may be used if they can be shown to lead to the same results. system

F.1 Reagents
F.2 Apparatus
F.3 Procedure

Annex G (informative) Generation of external control RNA (EC RNA) stocks

G.1 General

G.2 Reagents and apparatus
G.3 Linearization of plasmid DNA
G.4 In vitro RNA transcription
G.5 DNA contamination check
G.6 Quantification of EC RNA

Annex H (informative) Typical optical plate layout

Annex I (informative) Method validation studies and performance characteristics

Page count: 40