

ISO 22634-2:2019-12 (E)

Cigarettes - Determination of benzo[a]pyrene in cigarette mainstream smoke using GC/MS - Part 2: Method using cyclohexane as extraction solvent

Contents		Page
Foreword		iv
Introduction		v
1 Scope		1
2 Normative references		1
3 Terms and definitions		1
4 Principle		1
5 Apparatus		2
6 Reagents		2
7 Standards		3
7.1 General		3
7.2 Primary B[a]P-d12 stock solution: 100 µg/ml		3
7.3 Secondary B[a]P-d12 spiking solution: 40 ng/ml		3
7.4 Primary B[a]P stock solution: 100 µg/ml		3
7.5 Secondary B[a]P stock solution: 1 000 ng/ml		3
7.6 Working standard solutions		3
7.7 Storage of standard solutions		3
8 Preparation of sample		4
8.1 Sampling		4
8.2 Smoking		4
8.3 Glass-fibre filter pad extraction		4
8.4 Sample clean-up		4
8.5 Blank solution		5
9 Determination		5
9.1 GC/MS operating conditions		5
9.2 Calibration		6
9.3 Determination of B[a]P		6
9.4 Calculation		6
10 Repeatability and reproducibility		6
11 Test report		7
11.1 General		7
11.2 Characteristic data about the cigarette		7
11.3 Data about sampling		8
11.4 Description of the test		8
11.5 Test results		8
Annex A (informative) Example of a chromatogram of a cigarette smoke extract		9
Bibliography		10