ISO 21322:2020 (E)

Cosmetics — Microbiology — Testing of impregnated or coated wipes and masks

Contents

Foreword

Introduction

1 Scope

2 Normative references

3 Terms and definitions

4 Principle

4.1 General information
4.2 Selection of the test sample
4.3 Selection of the method
4.4 Recovery of microorganisms from the test sample
4.5 Enumeration of aerobic mesophilic microorganisms
4.5.1 General
4.5.2 Plate count method overview
4.5.3 Membrane filtration method overview
4.6 Detection of specified microorganisms by enrichment method

5 Diluents, neutralizers and culture media

5.1 General
5.2 Diluents and neutralizers
5.3 Culture media
5.3.1 Media for enumeration and detection
5.3.2 Media for preparation of spores of Bacillus subtilis

6 Apparatus and glassware

7 Strains of microorganisms

8 Handling of cosmetic products and laboratory samples

9 Procedure

9.1 General recommendation
9.2 Selection and preparation of the test sample
9.2.1 Selection of the test sample
9.2.2 Preparation of the initial suspension
9.3 Recovery of microorganisms
9.3.1 General
9.3.2 Stomaching
9.3.3 Shaking/Stirring
9.4 Enumeration of aerobic mesophilic microorganisms
9.4.1 General
9.4.2 Pour plate method
9.4.3 Surface spread method
9.4.4 Membrane filtration method
9.4.5 Incubation
9.4.6 Counting of colonies
9.5 Detection of specified microorganisms by enrichment method
9.5.1 General
9.5.2 Test for specified microorganisms

9.5.2.1 Preparation of the initial suspension in the enrichment broth
9.5.2.2 Incubation of the initial suspension
9.5.2.3 Isolation
9.5.2.4 Detection of specified microorganisms

10 Expression of results

10.1 Enumeration of aerobic mesophilic microorganisms
10.2 Detection of specified microorganisms

11 Suitability test

12 Test report

Annex A (normative) Guidance on methods for microbiological testing of impregnated or coated products — Wipes and masks

A.1 Selection of the test sample
A.2 Selection of the enumeration method for aerobic mesophilic microorganisms
A.3 Recovery of microorganisms
A.4 Summary of procedure for microbiological control of wipes and masks

Annex B (informative) Expression and interpretation of results

B.1 Counting and expression of results
B.2 Examples of expression of results — Plate count: Pour plate method
B.2.1 Treatment of the test sample
B.2.2 Single dose eye masks — Test sample 1 unit
B.2.3 Pack of wipes — Test sample 1/5 of a unit (test sample was a large towelette and therefore 1/5 of the towelette was weighed and tested)
B.2.4 Single dose face mask — Test sample 1/2 unit
B.2.5 Single dose eyes mask — Test sample 2 units (1 unit is less than 1 g)
B.2.6 Single dose face mask — Test sample 1 unit
B.3 Examples of expression of results — Plate count membrane filtration method
B.3.1 Treatment of the test sample
B.3.2 Pack of wipes — Test sample 1 unit
B.3.3 Single dose face mask — Test sample — 1/2 unit
B.3.4 Single dose eyes mask — Test sample 2 units — Each unit is less than 1 g
B.3.5 Single dose face mask — Test sample 1 unit
B.3.6 Single dose face mask — Test sample 1/2 unit

Annex C (normative) Suitability test method

C.1 Preparation of inoculum
C.1.1 Bacteria
C.1.2 Candida albicans
C.1.3 Preparation of spores of Bacillus subtilis
C.1.3.1 Culture media
C.1.3.1.1 Tryptic soy agar (TSA) or soybean casein digest agar (SCDA)
C.1.3.1.2 Peptone glucose agar with meat and yeast extract
C.1.3.1.3 Nutrient agar with 0,3 % of manganese (II) sulfate
C.1.3.1.4 Peptone and meat extract agar with sodium chloride
C.1.3.2 Preparation of spores stock suspension
C.2 Bioburden recovery
C.3 Neutralization of the antimicrobial properties of the product
C.3.1 General
C.3.2 Enumeration method
C.3.3 Enrichment method
C.4 Expression of results
C.4.1 Bioburden recovery
C.4.2 Neutralization of the antimicrobial properties of the product
C.4.2.1 Enumeration method
C.4.2.2 Enrichment method

Page count: 26