

ISO 23904:2020 (E)

Cigarettes — Determination of selected phenolic compounds in cigarette mainstream smoke with an intense smoking regime using HPLC-FLD

Contents

| | |
|---------|--|
| | Foreword |
| | Introduction |
| 1 | Scope |
| 2 | Normative references |
| 3 | Terms and definitions |
| 4 | Principle |
| 5 | Apparatus |
| 6 | Reagents |
| 7 | Preparation |
| 7.1 | General |
| 7.2 | Preparation of solutions — Acetic acid, with a volume fraction of 1 % solution |
| 7.3 | Preparation of standards |
| 7.3.1 | Primary phenolic compounds stock solutions |
| 7.3.2 | Secondary phenolic compounds stock solutions |
| 7.3.3 | Phenolic compounds working standards |
| 8 | Sampling |
| 9 | Tobacco product preparation |
| 10 | Sample generation — Smoking of cigarettes |
| 10.1 | General |
| 10.2 | Smoking machine setup |
| 10.3 | Smoking |
| 11 | Sample analysis |
| 11.1 | Preparation of sample |
| 11.2 | Determination |
| 11.2.1 | HPLC-FLD operating conditions |
| 11.2.2 | Calibration |
| 11.2.3 | Calculation |
| 12 | Repeatability and reproducibility |
| 12.1 | General |
| 12.2 | Results of the 2013 collaborative study |
| 13 | Report |
| Annex A | (informative) Examples of chromatograms |