ISO 23906-2:2023-02 (E)

Cigarettes - Determination of benzo[a]pyrene in cigarette mainstream smoke with an intense smoking regime using GC/MS - Part 2: Method using cyclohexane as extraction solvent

Contents

Page

Foreword iv

Introduction v

1 Scope 1

2 Normative references 1

3 Termsanddefinitions 1

4 Principles 1

5 Apparatus 2

6 Reagents 3

7 Standards 3
7.1 General 3
7.2 Primary B[a]P-d12 stock solution: 100 µg/ml 3
7.3 Secondary B[a]P-d12 spiking solution: 40 ng/ml 3
7.4 Primary B[a]P stock solution: 100 µg/ml 3
7.5 Secondary B[a]P stock solution: 1 000 ng/ml 3
7.6 Working standard solutions 3
7.7 Storage of standard solutions 3

8 Preparation of sample 4
8.1 Sampling 4
8.2 Smoking 4
8.3 Glass-fibre filter pad extraction 4
8.4 Sample clean-up 4
8.5 Blank solution 5

9 Determination 5
9.1 GC/MS operating conditions 5
9.2 Calibration 6
9.3 Determination of B[a]P 6
9.4 Calculation 6

10 Repeatability and reproducibility 6

11 Test report 7
11.1 General 7
11.2 Test results 7

Annex A (informative) Example of a chromatogram of a cigarette smoke extract 8

Bibliography 9