# ISO 21114:2026-10 (E)

## Nicotine pouches - Determination of nicotine content using liquid chromatography ultraviolet spectroscopy (LC-UV)

## Contents

Page


Foreword ..... iv

Introduction ..... v

1 Scope ..... 1

2 Normative references ..... 1

Terms and definitions

3 Termsanddefinitions ..... 1

4 Principle ..... 1

5 Apparatus ..... 1

6 Reagents ..... 3

7 Preparation ..... 3
7.1 Preparation of glassware ..... 3
7.2 Preparation of solutions ..... 3
7.2.1 Ammonium formate, 500 mmol/l (used for preparing extraction solution (7.2.2)) ..... 3
7.2.2 Extraction solution, 100 mmol/l ammonium formate ..... 3
7.2.3 Ammonium acetate stock buffer, 500 mmol/l (used for mobile phases) ..... 4
7.2.4 Mobile phase A, 10 mmol/l ammonium acetate buffer + 0,025 % ammonium hydroxide ..... 4
7.2.5 Mobile phase B, 10 mmol/l ammonium acetate buffer in acetonitrile + 0,025 % ammonium hydroxide ..... 4
7.2.6 Needle wash, 40:40:20:2 of water:methanol: 2-propanol:formic acid ..... 4
7.2.7 Seal wash, 90:10 of water:acetonitrile ..... 4
7.3 Preparation of standards ..... 4
7.3.1 Nicotine standard stock solution: approx. 25 mg/ml ..... 4
7.3.2 Nicotine calibration standards ..... 4

8 Sample preparation and extraction ..... 5
8.1 General ..... 5
8.2 Sample preparation ..... 5
8.3 Sample extraction ..... 5

9 Sample extract analysis ..... 6
9.1 General ..... 6
9.2 Suggested UPLC parameters ..... 6
9.3 Suggested HPLC parameters ..... 7
9.4 System suitability ..... 7
9.5 Calibration ..... 7

10 Repeatability and reproducibility ..... 8

11 Test report ..... 9

Annex A (informative) Examples of UPLC chromatograms ..... 10

Annex B (informative) Examples of HPLC chromatograms ..... 12

Bibliography ..... 14