ISO 20031:2020-02 (E)

Radiological protection - Monitoring and dosimetry for internal exposures due to wound contam ination with radionuclides

Contents

Page

Foreword v

Introduction vi

1 Scope 1

2 Normative references 1

3 Termsanddefinitions 1

4 Symbols and abbreviated terms 5
4.1 Symbols 5
4.2 Abbreviated terms 5

5 Purpose and need for special monitoring programmes for internal exposures due to wound contamination with radionuclides 5

6 General aspects of wound contamination 6
6.1 Introduction 6
6.2 Category of wound contaminants 6
6.3 Types of wounds and their specific retention of radionuclides 7

7 Monitoring programmes to assess contamination via a wound 7
7.1 Introduction 7
7.2 Main steps for the monitoring and dosimetry for internal exposures due to wound contamination with radionuclides 7
7.3 Collection of information to characterize the contaminated wound 8
7.3.1 General 8
7.3.2 Information concerning the type of wound 9
7.3.3 Information concerning the radioactive contaminant 9
7.4 In vivo wound measurements 9
7.5 Systemic activity monitoring 10

8 Performance criteria for radiobioassay measurements 11

9 Procedure for local and systemic dose assessment 11
9.1 Local (wound site) dose assessment 11
9.2 Systemic dose assessment 11
9.3 Impact of medical intervention on dose assessment 13
9.3.1 Local chelation therapy and/or the excision of contaminated tissue from the wound 13
9.3.2 Decorporation therapy 13
9.4 Software tools for bioassay data interpretation 13
9.5 Uncertainties 14
9.5.1 General 14
9.5.2 Uncertainties on local dose assessment 14
9.5.3 Uncertainties on internal dose assessment 14
9.6 Quality assurance 14

10 Recording 15
10.1 Recording in vivo measurement results 15
10.2 Recording in vitro radiobioassay and treatment waste results 15

11 Documentation of the dose assessment 16

12 Reporting 16

Annex A (informative) **Schematic representation of NCRP wound model, default parameters for retention equations and default transfer rates for the wound model for the various categories of radionuclides in wounds (adapted from NCRP report 156 (2007)[3])** **17**

Annex B (informative) **Types of wounds and their specific retention of radionuclides** **20**

Annex C (informative) **Example of a summary sheet that should follow the contaminated worker during his initial care** **23**

Annex D (informative) **Overview of typical methods used for in vitro bioassay measurements** **24**

Annex E (informative) **Equivalent dose rate in a contaminated wound ($mSv \cdot h^{-1} \cdot kBq^{-1}$) and equivalent dose rate received by the skin ($mSv \cdot h^{-1} \cdot kBq^{-1} \cdot cm^2$) for selected radionuclides** **25**

Annex F (informative) **Committed effective dose coefficients for intake of selected radionuclides via a contaminated wound for all wound model categories (adapted from Toohey RE et al., 2014[11])** **27**

Annex G (informative) **The IDEAS Guidelines[14] provide guidelines for the estimation of committed doses from incorporation monitoring data in case of wound** **30**

Bibliography **31**