ISO 1833-2:2020 (E)

Textiles — Quantitative chemical analysis — Part 2: Ternary fibre mixtures

Contents

Foreword

1 Scope

2 Normative references

3 Terms and definitions

4 Principle

5 Reagents and apparatus

6 Conditioning and testing atmosphere

7 Sampling and pre-treatment of laboratory test sample

8 Procedure

9 Calculation and expression of results

- 9.1 General
- 9.2 Calculation of percentages of mass of clean dry fibres, disregarding loss of fibre mass during pre-treatment
- 9.2.1 Variant 1
- 9.2.2 Variant 2
- 9.2.3 Variant 3
- 9.2.4 Variant 4
- 9.3 Calculation of the percentage of each component with adjustment by agreed moisture regains and, where appropriate, by correction factors for losses in mass during pre-treatment operations
- 9.4 Calculation of the quantitative analysis by manual separation
- 9.4.1 General
- 9.4.2 Calculation of the percentage mass of clean dry fibre disregarding loss in fibre mass during pre-treatment
- 9.4.3 Calculation of the percentage of each component with adjustment by agreed moisture regain and, where appropriate, by correction factors for losses in mass during pre-treatment

10 Method of quantitative analysis by a combination of manual separation and chemical means

11 Precision of methods

12 Test report

Annex A (informative) Examples of the calculation of percentages of the components of certain ternary mixtures using some of the variants described in 9.2

- A.1 Variant 1
- A.1.1 General
- A.1.2 Dry masses
- A.1.3 Masses after application of allowance for moisture
- A.2 Variant 4
- A.2.1 General
- A.2.2 Dry masses
- A.2.3 Mass after application of agreed moisture regains

Annex B (informative) Typical ternary mixtures which can be analysed using methods of analysis of binary mixtures specified in the parts of ISO 1833

Page count: 17