ISO 20136:2020 (E)

Leather — Determination of degradability by micro-organisms

Contents

Foreword

Introduction

1 Scope

2 Normative references

3 Terms and definitions

4 Symbols and abbreviated terms

5 Principle

5.1 General
5.2 Assessment of biodegradation by manual titration; method A
5.3 Assessment of biodegradation by infrared (IR) detection; method B

6 Chemicals

7 Apparatus and materials

8 Procedure

8.1 Collection and preparation of the inoculum
8.2 Preparation of the test material and reference material
8.3 Test conditions and incubation period
8.4 Termination of the test

9 Quantification

9.1 Assessment of biodegradation by manual titration (method A)
9.1.1 Determination of the organic carbon content
9.1.2 Determination of the amount of CO2 produced
9.1.3 Correcting for normality of HCl
9.1.4 Percentage of biodegradation from CO2 evolved
9.2 Assessment of biodegradation by IR (method B)
9.2.1 Determination of the organic carbon content
9.2.2 Determination of the amount of CO2 produced
9.2.3 Percentage of biodegradation from CO2 data
9.2.3.1 General
9.2.3.2 Expression of the total number of moles of CO2 evolved
9.2.3.3 Conversion of moles of CO2 evolved into mg of CO2 evolved
9.2.3.4 Calculation of the CO2 mass present in the initial test sample
9.2.3.5 Calculation of the CO2 mass evolved from the initial sample
9.2.3.6 Calculation of the percentage of biodegradation
9.2.3.7 Calculation sequence for the determination of sample biodegradation

10 Expression of results

11 Validity of results

12 Test report

Annex A (informative) Determination of the degree and rate of degradation of the material

Annex B (informative) Quantitative determination of leather biodegradation

Annex C (informative) Comparative biodegradability using different waste waters