ISO 20136:2020 (E)

Leather — Determination of degradability by micro-organisms

Contents

Foreword

Introduction

- 1 Scope
- 2 Normative references
- 3 Terms and definitions
- 4 Symbols and abbreviated terms
- 5 Principle
 - 5.1 General
 - 5.2 Assessment of biodegradation by manual titration; method A
 - 5.3 Assessment of biodegradation by infrared (IR) detection; method B
- 6 Chemicals
- 7 Apparatus and materials

8 Procedure

- 8.1 Collection and preparation of the inoculum
- 8.2 Preparation of the test material and reference material
- 8.3 Test conditions and incubation period
- 8.4 Termination of the test
- 9 Quantification
 - 9.1 Assessment of biodegradation by manual titration (method A)
 - 9.1.1 Determination of the organic carbon content
 - 9.1.2 Determination of the amount of CO2 produced
 - 9.1.3 Correcting for normality of HCI
 - 9.1.4 Percentage of biodegradation from CO2 evolved
 - 9.2 Assessment of biodegradation by IR (method B)
 - 9.2.1 Determination of the organic carbon content
 - 9.2.2 Determination of the amount of CO2 produced
 - 9.2.3 Percentage of biodegradation from CO2 data
 - 9.2.3.1 General
 - 9.2.3.2 Expression of the total number of moles of CO2 evolved
 - 9.2.3.3 Conversion of moles of CO2 evolved into mg of CO2 evolved
 - 9.2.3.4 Calculation of the CO2 mass present in the initial test sample
 - 9.2.3.5 Calculation of the CO2 mass evolved from the initial sample
 - 9.2.3.6 Calculation of the percentage of biodegradation
 - 9.2.3.7 Calculation sequence for the determination of sample biodegradation
- 10 Expression of results
- 11 Validity of results
- 12 Test report
- Annex A (informative) Determination of the degree and rate of degradation of the material
- Annex B (informative) Quantitative determination of leather biodegradation
- Annex C (informative) Comparative biodegradability using different waste waters